

1 CLAIMS

2

3 1. A method of preparing a composition for the
4 treatment of dermatological disorders, wherein coal
5 tar is filtered to remove impurities by filter
6 compression.

7

8 2. A method as in Claim, wherein zinc pyrithione is
9 incorporated into the composition.

10

11 3. A method as in any of the previous Claims, wherein
12 the coal tar is filtered by being fed through a
13 compressed charcoal filter.

14

15 4. A method as in Claim 3, wherein the coal tar is fed
16 through the filter at 17 to 19 psi.

17

18 5. A method as in Claim 3, wherein the coal tar is fed
19 through the filter at 18 psi.

20

21 6. A method as in Claims 3 to 5, wherein compressed air
22 is used to force the coal tar through the filter.

23

24 7. A method as in Claims 3 to 6, wherein the coal tar
25 is left in the filter system for 8 hours.

26

27 8. A method as in Claims 3 to 7, wherein the fluid that
28 has been passed through the filter is boiled.

29

30 9. A method as in Claim 8, wherein the fluid that has
31 been passed through the filter is boiled for 5
32 minutes.

33

- 1 10. A method as in Claims 8 or 9, wherein the boiled,
2 filtered fluid is allowed to cool to room
3 temperature.
4
- 5 11. A method as in Claims 8, 9 or 10, wherein the top
6 layer of the boiled, filtered fluid is refiltered.
7
- 8 12. A method as in Claim 11, wherein refiltering is
9 through a nylon mesh.
10
- 11 13. A method as in any of the previous Claims, where a
12 surfactant is added to the formulation.
13
- 14 14. A method as in Claim 13, wherein the surfactant is
15 an ionic surfactant.
16
- 17 15. A method as in Claims 13 or 14, wherein the
18 surfactant is sodium lauryl sulphate.
19
- 20 16. A method as in any of the previous Claims, wherein a
21 carrier is added to the formulation.
22
- 23 17. A method as in any of the previous Claims, wherein
24 multiple carriers are added to the formulation.
25
- 26 18. A method as in Claims 16 or 17, wherein the carrier
27 is isopropyl myristate.
28
- 29 19. A method as in Claims 16 or 17, wherein the carrier
30 is ethyl alcohol.
31

1 20. A method as in any of the previous Claims, wherein
2 the formulation is placed in a spray or aerosol
3 container.
4

5 21. A composition for the treatment of dermatological
6 disorders comprising:
7

- 8 • coal tar;
- 9 • zinc pyrithione;
- 10 • one or more surfactants; and
- 11 • one or more carriers

12

13 22. A method as in Claim 21, wherein the composition
14 also contains allantoin.
15

16 23. A composition as in Claims 21 and 22, wherein the
17 surfactant is an ionic surfactant.
18

19 24. A composition as in Claims 21 to 23, wherein the
20 surfactant is sodium lauryl sulphate.
21

22 25. A composition as in Claims 21 to 24, wherein the
23 carrier is isopropyl myristate.
24

25 26. A composition as in Claims 21 to 24, wherein the
26 carrier is ethyl alcohol.
27

28 27. A composition as in Claims 21 to 25, wherein the
29 composition also comprises an anti-fungal agent.
30

31 28. A composition as in Claim 27, wherein the anti-
32 fungal agent is undecylenic acid.
33

1 29. A composition for the treatment of dermatological
2 disorders, comprising:

3

- 4 • zinc pyrithione;
- 5 • alcoholic extract of coal tar;
- 6 • allantoin;
- 7 • sodium lauryl sulphate;
- 8 • isopropyl myristate;
- 9 • ethyl alcohol; and
- 10 • undecylenic acid

11

12 30. A composition as in Claim 29, wherein the
13 ingredients in the following amounts:

14

- | | |
|------------------------------------|--------|
| 15 • zinc pyrithione | 0.20% |
| 16 • alcoholic extract of coal tar | 0.25% |
| 17 • allantoin | 0.25% |
| 18 • sodium lauryl sulphate | 0.10% |
| 19 • isopropyl myristate | 49.45% |
| 20 • ethyl alcohol | 49.45% |
| 21 • undecylenic acid | 0.30% |

22

23 31. A composition as in Claims 21 to 30, wherein the
24 composition is provided in a spray form.

25

26 32. A composition as in Claims 21 to 30, wherein the
27 composition is provided in an aerosol form.

28